

Nonlinear mixed Cherepanov boundary-value problem

Obnosov Y., Zulkarnyaev A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Informa UK Limited, trading as Taylor & Francis Group We consider a nonlinear boundary-value problem consisting in the determination of a function (Formula presented.) that is meromorphic in the upper half-plane, satisfies the homogeneous Hilbert boundary condition on the set L of n intervals of the real axis and has a given module on the set (Formula presented.). This problem was stated and solved in Cherepanov. Cherepanov proved that the required solution with a given number and location of its interior zeros and poles and with integrable singularities at all endpoints of L exists if and only if $n-1$ solvability conditions are satisfied. We prove that this problem is unconditionally solvable. A particular solution, (Formula presented.), is found in the class of meromorphic functions with a properly chosen location of $n-1$ zeros and poles. Namely, we have shown that these zeros and poles are defined as the solution of some solvable real analogue of the Jacobi inversion problem. A general meromorphic solution of the Cherepanov problem is obtained with the help of the particular solution (Formula presented.). The problem of a possible decrease in the number of zeros and poles of the desired solution is investigated.

<http://dx.doi.org/10.1080/17476933.2018.1493465>

Keywords

analytic functions, closed form solution, Nonlinear mixed boundary-value problem